


Transdermal Estrogen

Not intended to diagnose, treat, prevent, or cure any disease.



Contains estradiol (CAS #50-28-2), ethanol, propylene glycol USP, diethylene glycol monoethyl ester USP, distilled water. Total estradiol 0.24% by volume.


Take 0.57mL (approximately 1/4 dropperful) per day applied to neolabia or scrotum topically for standard feminizing effects. Do not inject, eat, boof, or otherwise ingest internally. For lesser reduced effects apply to forearm or underarm. Keep dry for one hour after application.

Recommend researching potential side effects and complications of hormone replacement therapy before use – <https://transfemscience.org> is a great place to start!

Email mablesbakerymutualaid@proton.me with any questions, to report any issues or to report your results.

Transdermal Estrogen

Not intended to diagnose, treat, prevent, or cure any disease.



Contains estradiol (CAS #50-28-2), ethanol, propylene glycol USP, diethylene glycol monoethyl ester USP, distilled water. Total estradiol 0.24% by volume.


Take 0.57mL (approximately 1/4 dropperful) per day applied to neolabia or scrotum topically for standard feminizing effects. Do not inject, eat, boof, or otherwise ingest internally. For lesser reduced effects apply to forearm or underarm. Keep dry for one hour after application.

Recommend researching potential side effects and complications of hormone replacement therapy before use – <https://transfemscience.org> is a great place to start!

Email mablesbakerymutualaid@proton.me with any questions, to report any issues or to report your results.

Transdermal Estrogen

Not intended to diagnose, treat, prevent, or cure any disease.



Contains estradiol (CAS #50-28-2), ethanol, propylene glycol USP, diethylene glycol monoethyl ester USP, distilled water. Total estradiol 0.24% by volume.


Take 0.57mL (approximately 1/4 dropperful) per day applied to neolabia or scrotum topically for standard feminizing effects. Do not inject, eat, boof, or otherwise ingest internally. For lesser reduced effects apply to forearm or underarm. Keep dry for one hour after application.

Recommend researching potential side effects and complications of hormone replacement therapy before use – <https://transfemscience.org> is a great place to start!

Email mablesbakerymutualaid@proton.me with any questions, to report any issues or to report your results.

Transdermal Estrogen

Not intended to diagnose, treat, prevent, or cure any disease.



Contains estradiol (CAS #50-28-2), ethanol, propylene glycol USP, diethylene glycol monoethyl ester USP, distilled water. Total estradiol 0.24% by volume.

Take 0.57mL (approximately 1/4 dropperful) per day applied to neolabia or scrotum topically for standard feminizing effects. Do not inject, eat, boof, or otherwise ingest internally. For lesser reduced effects apply to forearm or underarm. Keep dry for one hour after application.

Recommend researching potential side effects and complications of hormone replacement therapy before use – <https://transfemscience.org> is a great place to start!

Email mablesbakerymutualaid@proton.me with any questions, to report any issues or to report your results.